

Topological rigidity of manifolds and the Farrell—Jones Conjecture

Guglielmo Nocera

October 20th, 2026

Contents

1	K_1 and h-cobordisms	1
2	The Borel Conjecture	2
3	K-theory and the Farrell-Jones conjecture	2
4	Categorification	3

1 K_1 and h-cobordisms

Recall the projective class group/Grothendieck group of a ring

$$K_0(R) = \left(\bigoplus_{P \text{ f.g. proj } R\text{-module}} \mathbb{Z} \langle P \rangle \right) / \text{short exact sequence.}$$

Definition 1.1. Let R be a (not necessarily commutative) ring. Let $GL(R) = \bigcup_n GL_n(R)$ (as a group, where each one is embedded into the other by completing with an identity block). We define

$$K_1(R) := GL(R)^{\text{ab}} = GL(R) / [GL(R), GL(R)] \simeq GL(R) / \{\text{elementary matrices}\}.$$

Definition 1.2. Let G be a group. Consider the homomorphism

$$\begin{aligned} f : G \times \{\pm 1\} &\rightarrow K_1(\mathbb{Z}[G]) \\ (g, \pm 1) &\mapsto \pm g \in GL(\mathbb{Z}[G]) \end{aligned}$$

The Whitehead group of G is

$$\text{Wh}(G) = \text{coker}(f) = GL(\mathbb{Z}[G]) / (G \times \{\pm 1\} \times \{\text{elementary matrices}\}).$$

Definition 1.3. Let M be a topological n -manifold. An h-cobordism over M is an $n + 1$ -manifold W with boundary such that

- $\partial W \simeq M \amalg N$ where N is an n -manifold
- $M \hookrightarrow W, N \hookrightarrow W$ are homotopy equivalences.

We denote

$$\mathcal{H}(M) = \{\text{h-cobordisms over } M\} / \text{homeomorphisms rel } M.$$

DRAW PICTURE

Theorem 1.4 (s-cobordism). *If M is a closed manifold of dimension ≥ 5 , there is a bijection*

$$\mathcal{H}(M) \leftrightarrow \text{Wh}(\pi_1(M)).$$

Corollary 1.5. *Let G be a discrete group. If $\text{Wh}(G) = 0$, then for every closed manifold M of dimension ≥ 5 and $\pi_1(M) = G$, every h -cobordism over M is trivial.*

2 The Borel Conjecture

Definition 2.1. A closed manifold N is called topologically rigid if any homotopy equivalence $M \rightarrow N$ with a closed manifold M as source is homotopic to a homeomorphism.

COUNTEREXAMPLE IN NONCLOSED CASES: ANNULUS AND CIRCLE

Definition 2.2. A topological space X is aspherical if $\pi_i(X) = 0$ for all $i \neq 0$.

Conjecture 2.3 (Borel Conjecture for G). *Let G be a discrete finitely presented torsionfree group. Any closed aspherical manifold M with $\pi_1(M) \simeq G$ is topologically rigid.*

The conjecture is true in dimension 1 and 2 by classification, in dimension 3 as a consequence of Perelman's proof of Thurston's Geometrization Conjectures, and open in dimension 4.

It is also open dimension ≥ 5 , but proven for many interesting groups by using a K-theoretic criterion, in the same spirit as the vanishing of $\text{Wh}(\pi_1(M))$ implies cobordism-rigidity.

3 K-theory and the Farrell-Jones conjecture

Recall 3.1. There exists a whole algebraic K-theory spectrum of a ring R , which we denote by $K_*(R)$.

$$\pi_i(K_*(R)) = K_i(R)$$

(this is a theorem for 0, 1 and a definition for ≥ 2).

There exists also something called L-theory, which we will skip. Instead of parametrizing modules up to short exact sequence, automorphisms, etc., it parametrizes modules with quadratic/symmetric forms up to Lagrangian, etc.

Conjecture 3.2 (K-theoretic Farrell-Jones, simplified). *Slogan: “K-theory of integral group rings is a homology theory!”*

*Let G be a finitely presented torsionfree group. Then the **assembly map***

$$\text{asmb}_K : \Sigma_+^\infty BG \wedge K_*(\mathbb{Z}) \rightarrow K_*(\mathbb{Z}[G])$$

is an equivalence/isomorphism of spectra.

Note: the assembly map exists, the conjecture only says that it is an equivalence.

Theorem 3.3 (Farrell–Jones). *The K-theoretic Farrell–Jones Conjecture, combined with the analogous statement for L-theory, implies the Borel Conjecture in $\dim \geq 5$.*

The Farrell–Jones Conjecture has been proven in various cases in works by Lück, Bartels, Reich, et al.: true for

- hyperbolic groups
- finite-dimensional CAT(0) groups
- virtually solvable groups, etc.
- fundamental groups of connected manifolds of dimension ≤ 3
- certain lattices
- $\mathrm{GL}_n(\mathbb{Q}), \mathrm{GL}_n(\mathbb{F}_p(t))$
- $\Gamma_{g,r}^s$, the mapping class group of a closed orientable surface of genus g, r boundary components and s punctures

Also, they show that the class for which the Conjecture is true is closed under:

- subgroups
- overgroups of fte index
- finite direct products,
- directed colimits
- certain extensions
- all free products.

The fact that it is closed under directed colimits makes it harder to build counterexamples.

4 Categorification

K_* is an invariant of categories:

$$R \rightsquigarrow \mathrm{Proj}_R \rightsquigarrow K(\mathrm{Proj}_R) \simeq K_*(R)$$

but this can be generalized to stable ∞ -categories, that is:

Theorem 4.1. *There is a functor $K : \{\text{stable } \infty\text{-categories}\} \rightarrow \{\text{spectra}\}$ s.t. $K(\mathrm{Perf}_R) = K_*(R)$, sending “fiber-cofiber sequences” to short exact sequences.*

Goal 4.2. Build a fiber-cofiber sequence of stable ∞ -categories

$$\mathcal{N} \xrightarrow{\Phi} \mathcal{C} \xrightarrow{\Psi} \mathcal{D}$$

such that $K(\Phi) = \mathrm{asmb}_K$. Give manageable conditions for which $K(\mathcal{N}) = 0$ as a spectrum, and hence asmb_K is an equivalence.

The answer turns out to involve a more powerful invariant, namely continuous K-theory. In work with Levin and Saunier, we achieved the goal for the L-theory version.